
-continued

<223> OTHER INFORMATION: Primer DNA sequence of Beta-actin-R

<400> SEQUENCE: 8

agctcgtagc ttttctcca

19

<210> SEQ ID NO 9

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer DNA sequence of GAPDH-F

<400> SEQUENCE: 9

atgacatcaa gaagggtggtg aa

22

<210> SEQ ID NO 10

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer DNA sequence of GAPDH-R

<400> SEQUENCE: 10

gctgttgaag tcagaggaga c

21

<210> SEQ ID NO 11

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Guide RNA sequence of CXCR4 gRNA no 1

<400> SEQUENCE: 11

tgttggctgc cttaactacat

20

<210> SEQ ID NO 12

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Guide RNA sequence of HRH gRNA no 3

<400> SEQUENCE: 12

cgtatcaagtc cgccaccgag

20

<210> SEQ ID NO 13

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Guide RNA sequence of non-targeting gRNA

<400> SEQUENCE: 13

acggaggcta agcgctcgcaa

20

1-173. (canceled)

174. A method of mobilizing a stem cell in a subject, the method comprising:

administering to the subject an inhibitor of CXCR4 and an inhibitor of GPCR_x;

wherein the combination of the inhibitor of CXCR4 and the inhibitor of GPCR_x induces an enhanced amount of

stem cell mobilization relative to the amount of stem cell mobilization induced by the CXCR4 inhibitor only.

175. A method of mobilizing a stem cell in a subject having a CXCR4 protomer and a GPCR_x protomer, the method comprising:

administering to the subject an inhibitor of CXCR4 and an inhibitor of GPCR_x;